Information Dissemination in Schools: The Facts about Adolescent Suicide


ORANGE COUNTY • NEW YORK

This document is a true and false test on adolescent suicide, which could be presented to staff as well as parents as a way of increasing their awareness and knowledge. By simply giving this true and false to staff and parents and allowing for some time to discuss questions and concerns, schools can effectively increase awareness about adolescent suicide and may help prevent an incident of suicide in their school.

True/False Test

True False

	1.	Adolescent suicide is an increasing problem in the United States.
		Most teenagers will reveal that they are suicidal or have emotional problems for which they would like emotional help.
	3.	Adolescents who talk about suicide do not attempt or kill themselves.
		Educating teens about suicide leads to increased suicide attempts, since it provides them with ideas and methods about killing themselves.
		Talking about suicide in the classroom will promote suicidal ideas and suicidal behavior.
	6.	Parents are often unaware of their child's suicidal behavior.
		The majority of adolescent suicides occur unexpectedly without warning signs.
	8.	Most adolescents who attempt suicide fully intend to die.
		There is a significant difference between male and female adolescents regarding suicidal behavior.
		The most common method for adolescent suicide deaths is drug overdose.
		Because female adolescents die by suicide at a lower rate than male adolescents, their attempts should not be taken seriously.
	12.	Not all adolescents who engage in suicidal behavior are mentally ill.
	13.	Suicidal behavior is inherited.
	14.	Adolescent suicide occurs only among poor adolescents.
		The only one who can help a suicidal adolescent is a counselor or a mental health professional.
		Adolescents cannot relate to a person who has experienced suicidal thoughts.
		If an adolescent wants to kill him/herself, there is nothing anyone can do to prevent its occurrence.



True/False



Orange-Ulster BOCES 845 291-0100 www.ouboces.org



Orange County Department of Mental Health 845 291-2600

www.orangecountygov.com

Suggested Citation: Roggenbaum, S., Doan, J., & Lazear, K.J. (2014). Youth suicide prevention school-based guide: Orange County, New York—True/False 1: Information dissemination in schools—The facts about adolescent suicide. Tampa, FL: University of South Florida, College of Behavioral and Community Sciences, Louis de la Parte Florida Mental Health Institute (FMHI Series Publication #256-1t-rev).

This publication is also available on-line as an Adobe Acrobat PDF file: <u>http://www.ouboces.org</u> and <u>http://www.orangecountygov.com</u> Select County Departments, Department of Mental Health

- Adolescent suicide has been increasing dramatically in the United States. False. While one suicide is one too many, the youth suicide rate has been declining over the past two decades. Other than a couple of minor increases in 1994 (1%) and in 2000 (3%), the child and adolescent suicide rate consistently declined for 15 years, deceasing from 4.4/100,000 in 1988 to 2.83/100,000 in 2003. The youth suicide rate for 10-24 year olds rose 8% from 2003 to 2004, then showed a general decline through 2007 but the rate increased again in 2008 by 4.5%, the most current year data is available as of this publication (2). The 2008 suicide rate for 15–19 year olds stands at 7.42 per 100,000 and the 2008 suicide rate for 10–14 year olds is 1.07 per 100,000 (2).
- 2. Most teenagers will reveal that they are suicidal or have emotional problems for which they would like emotional help. True. Most teens will reveal that they are suicidal and although studies have shown that they are more willing to discuss suicidal thoughts with a peer than a school staff member, this disposition that most teens have towards expressing suicidal ideations could be used for screening adolescents through questionnaires and/or interviews (4).
- 3. Adolescents who talk about suicide do not attempt suicide or kill themselves. False. One of the most ominous warning signs of adolescent suicide is talking repeatedly about one's own death. Adolescents who make threats of suicide should be taken seriously and provided the help that they need. In this manner, suicide attempts can be averted and lives can be saved (3, 6).
- 4. Educating teens about suicide leads to increased suicide attempts, since it provides them with ideas and methods about killing themselves. False. When issues concerning suicide are taught in a sensitive, educational context they do not lead to, or cause, further suicidal behaviors. Since three-fourths (77%) of teenage students state that if they were contemplating suicide they would first turn to a friend for help, peer assistance programs have been implemented throughout the nation. These educational programs help students to identify peers at risk and help them receive the help they need. Such programs have been associated with increased student knowledge about suicide warning signs and how to contact a hotline or crisis center, as well as increased likelihood to refer other students at risk to school counselors and mental health professionals. Furthermore, directly asking an adolescent if he or she is thinking about suicide displays

care and concern and may aid in clearly determining whether or not an adolescent is considering suicide. Research shows that when issues concerning suicide are taught in a sensitive and educational manner, students demonstrate significant gains in knowledge about the warning signs of suicide and develop more positive attitudes toward help- seeking behaviors with troubled teens (1, 7, 8, 11, 27).

- 5. Talking about suicide in the classroom will promote suicidal ideas and suicidal behavior. False. Talking about suicide in the classroom provides adolescents with an avenue to talk about their feelings, thereby enabling them to be more comfortable with expressing suicidal thoughts and increasing their chances of seeking help from a friend or school staff member. Additionally, recent research indicated that asking about suicidal behavioral does not plant the idea of suicidal ideation or behavior in a screening survey were no more likely to report thinking about suicide than students not exposed to these questions. Also, the research results seem to indicate that asking about suicidal ideation or behavior may have been helpful for at-risk students (i.e., those with depression symptoms or previous suicide attempts) (3, 27, 28).
- 6. Parents are often unaware of their child's suicidal behavior. True. One study has shown that as much as 86% of parents were unaware of their child's suicidal behavior. Another study found that parents were unaware of their children's depressive symptoms, as well as their alcohol use, both risk factors for youth suicidal behavior (1, 3, 30).
- 7. The majority of adolescent suicides occur unexpectedly without warning signs. False. Nine out of ten adolescents who die by suicide give clues to others before their suicide attempt. Warning signs for adolescent suicide include depressed mood, substance abuse, loss of interest in once pleasurable activities, decreased activity levels, decreased attention, distractability, isolation, withdrawing from others, sleep changes, appetite changes, morbid ideation, offering verbal cues (i.e., "I wish I were dead"), offering written cues (i.e., notes, poems), and giving possessions away. In addition, the following risk factors place an adolescent at increased risk for suicidal behavior: having a previous suicide attempt, having a recent relationship breakup, being impulsive, having low self-esteem, being homosexual, coming from an abusive home, having easy access to a firearm, having low grades, and being exposed to suicide or suicidal behavior by another person. Moreover, most suicidal

Answers to True/False Test continued

adolescents attempt to communicate their suicidal thoughts to another in some manner. Not surprisingly, an effective way to prevent adolescent suicide is to learn to identify the warning signs that someone is at risk (4, 5, 9, 11, 15-18, 29-34).

- 8. Most adolescents who attempt suicide fully intend to die. False. Most suicidal adolescents do not want suicide to happen. Rather, they are torn between wanting to end their psychological pain through death and wanting to continue living, though only in a more hopeful environment. Such ambivalence is communicated to others through verbal statements and behavior changes in 80% of suicidal youths. (1, 10).
- **9.** There is a significant difference between male and female adolescents regarding suicidal behavior. True. Adolescent females are significantly more likely than adolescent males to have thought about suicide and to have attempted suicide. More specifically, adolescent females are 1.5 to 2 times more likely than adolescent males to report experiencing suicidal ideation and 3 to 4 times more likely to attempt suicide. Adolescent males are 4 to 5.5 times more likely than adolescent females attempt. While adolescent females to complete a suicide attempts, adolescent males time out of 25 suicide attempts, adolescent males kill themselves in one out of every three attempts (1, 3, 4, 7, 12).
- 10. The most common method for adolescent suicide death is drug overdose. False. In 2007, young people were much more likely to use firearms, suffocation, and poisoning than other methods of suicide, overall. However, while adolescents (ages 15-19) were more likely to use firearms than suffocation, children (ages 10-14) were dramatically more likely to use suffocation. Having a gun in the house increases an adolescent's risk of suicide. Regardless of whether a gun is locked up or not, its presence in the home is associated with a higher risk for adolescent suicide. This is true even after controlling for most psychiatric variables. Homes with guns are 4.8 times more likely to experience a suicide of a resident than homes without guns. In lieu of these findings, it should not be surprising that restricting access to handguns has been found to significantly decrease suicide rates among 15-24 year olds (1, 3, 12, 13, 15, 23, 24, 25).

- 11. Because female adolescents die by suicide at a lower rate than male adolescents, their attempts should not be taken seriously. False. One of the most powerful predictors of death by suicide is a prior suicide attempt. Adolescents who have attempted suicide are 8 times more likely than adolescents who have not attempted suicide to attempt suicide again. Between one-third to one-half of adolescents who kill themselves have a history of a previous suicide attempt. Therefore, all suicide attempts should be treated seriously, regardless of sex of the attempter (1, 3, 4, 12, 15, 16-21).
- **12.** Not all adolescents who engage in suicidal behavior are mentally ill. True. The majority of adolescents have entertained thoughts about suicide at least once in their lives. There are cases of some adolescents attempting and dying by suicide who do not appear to have a diagnosable mental disorder. However, research studies regarding adolescents who die by suicide suggest that most (evidence suggests over 90%) have a diagnosable, although not always diagnosed, mental health disorder at the time of their death. Additionally, research suggests that identifying at-risk youth, by utilizing depression scales and psychopathology inventories, through screening and treating those individuals who test positive for mental illness can benefit from counseling by a trained professional (11, 14).
- 13. Suicidal behavior is inherited. False. There is no specific suicide gene that has ever been identified. Studies involving twins have found higher concordance rates for suicide in monozygotic twins than in dizygotic twins; meaning that an identical twin would be more likely than a fraternal twin to engage in suicidal behavior if his/her co-twin died by suicide. However, no study to date has examined the concordance for suicide in monozygotic twins separated at birth and raised apart, a requirement necessary to be met as a means to indicate inheritance of psychiatric illness. Such a study could assess the effects that parental rearing style and familial environment have on suicidal behavior. Interestingly enough, when compared to control subjects, adolescent suicide victims have been found to have had significantly less frequent and less satisfying communication with their parents (1, 12, 20, 21).

Answers to True/False Test continued

- 14. Adolescent suicide occurs only among poor adolescents. False. Adolescent suicide occurs in all socioeconomic groups. Socioeconomic variables have not been found to be reliable predictors of adolescent suicidal behavior. Instead of assessing adolescents'socioeconomic backgrounds, school professionals should assess their social and emotional characteristics (i.e., affect, mood, social involvement, etc.) to determine if they are at increased risk (1, 3, 15, 16, 21).
- **15. The only one who can help a suicidal adolescent is a counselor or a mental health professional. False.** Most adolescents who are contemplating suicide are not presently seeing a mental health professional. Rather, most are likely to approach a family member, peer, or school professional for help. Displaying concern and care as well as ensuring that the adolescent is referred to a mental health professional are ways paraprofessionals can help (7).
- **16. Adolescents cannot relate to a person who has experienced suicidal thoughts. False.** Data from the 2009 Youth Risk Behavior Surveillance Survey (YRBSS), which surveyed 16,220 high school students, found that more than one in ten students (13.8%) had seriously considered attempting suicide in the previous year. A population study of 5,000 teenagers from a rural community showed that 40% had entertained ideas of suicide at some point in their lives. Some researchers have estimated that it is more realistic that greater than half of all high school students have experienced thoughts of suicide (1, 14, 22).
- 17. If an adolescent wants to kill him/herself, there is nothing anyone can do to prevent its occurrence. False. One of the most important things an individual can do to prevent suicide is to identify the warning signs of suicide and recognize an adolescent at increased risk for suicide. School professionals should, therefore, be aware of these risk factors and know how to respond when a student threatens or attempts suicide. The existence of a school crisis intervention team may assist with this process (3, 11, 14, 26).

References

Information Dissemination in Schools

- 1. King, K.A. (1999). Fifteen prevalent myths about adolescent suicide. *Journal of School Health, 69*(4), 159–161.
- Centers for Disease Control and Prevention: Web-based Injury and Statistics Query and Reporting System. (2011). *Fatal injury reports, National and Regional, 1999-2008.* Retrieved from http://webappa.cdc.gov/sasweb/ncipc/ mortrate10_sy.html
- 3. Zenere, F.J., & Lazarus, P.J. (1997). The decline of youth suicidal behavior in an urban, multicultural public school system following the introduction of a suicide prevention and intervention program. *Suicide and Life-Threatening Behavior*, *27*(4), 387–403.
- 4. Gould, M.S., & Kramer, R.A. (2001). Youth suicide prevention. *Suicide and Life-Threatening Behavior, 31*, 6–31.
- Dube, S. R., Anda, R. F., Felitti, V. J., Chapman, D. P., Williamson, D., & Giles, W.H. (2001). Childhood abuse, household dysfunction, and the risk of attempted suicide throughout the lifespan: Findings from the Adverse Childhood Experiences Study. JAMA, 286(24), 3089-3096.
- 6. Kirk, W.G. (1993). *Adolescent suicide: A school-based approach to assessment and intervention*. Champain, IL: Research Press.
- 7. Tierney, R., Ramsay, R., Tanney, B., & Lang, W. (1991). Comprehensive school suicide prevention programs. In A.A. Leenaars & S. Wenkstern (Eds.), *Suicide Prevention in Schools*. New York: Hemishere Publishing Corporation.
- 8. Kalafat, J., & Elias, M. (1994). An evaluation of a schoolbased suicide awareness intervention. *Suicide and Life-Threatening Behavior, 24*(3), 224–233.
- 9. Haas, A.P., Eliason, M., Mays, V.M., Mathym R.M., Cochran, S.D., D'Augelli, A.R. . . . Clayton, P.J. (2001). Suicide and suicide risk in lesbian, gay, bisexual, and transgender populations: Review and recommendations. *Journal of Homosexuality, 58*(1), 10-51.
- King, K.A., Price, J.H., Telljohann, S.K., & Wahl, J. (1999). High school health teachers' knowledge of adolescent suicide. *American Journal of Health Studies*, *15*(3), 156–163.
- 11. Gould, M., Greenberg, T., Velting, D., & Shaffer, D. (2003). Youth suicide risk and preventive interventions: A review of the past 10 years. Journal of the *American Academy of Child and Adolescent Psychiatry*, *42*(4), 386–405.
- O'Carroll, P.W., Potter, L.B., & Mercy, J.A. (1994). Programs for the prevention of suicide among adolescents and young adults. *MMWR CDC Surveillance Summary 43* (RR-6). 1–7.

- Shaffer, D., Garland, A., Vieland, V., Underwood, M., & Busner, C. (1991). The impact of curriculum-based suicide prevention programs for teenagers. *Journal of the American Academy of Child and Adolescent Psychiatry*, 30(4), 588–596.
- 14. Garland, A.F., & Zigler, E. (1993). Adolescent suicide prevention: Current research and social policy implications. *American Psychologist*, *48*(2), 169–182.
- 15. Borowsky, I.W., Ireland, M., & Resnick, M.D. (2001) Adolescent suicide attempts: Risks and protectors. *Pediatrics*, *107*(3), 485–493.
- Moscicki, E. (1999). Epidemiology of Suicide. In D.G. Jacobs (ed), *The Harvard Medical School Guide to Suicide Assessment and Intervention*. (pp. 40–51). San Francisco: Jossey-Bass Publishing.
- 17. Shaffer D., Gould, M., & Fisher, P. (1996). Psychiatric diagnosis in child and adolescent suicide. *Archives of General Psychiatry*, *53*, 339–348.
- Shaffer, D., Pfeffer, C.R., & Work Group on Quality Issues. (2001). Practice parameter for the assessment and treatment of children and adolescents with suicidal behavior. *Journal of American Academy of Child and Adolescent Psychiatry*, 40(1) supp, 24–51.
- 19. Zametkin, A.J. Alter, M.R., & Yemini, T. (2001). Suicide in teenagers: Assessment, management, and prevention. *Journal of American Medical Association*, *286*(24). 3120–3125.
- 20. Lester, D. (2000). *Suicide prevention: Resources for the millennium*. Ann Arbor, MI: Sheridan Books.
- 21. Brent, D.A. (1995). Risk factors for adolescent suicide and suicidal behavior: Mental and substance abuse disorders, family environmental factors and life stress. *Suicide and Life-Threatening Behavior, 25,* 52–63.
- 22. Centers for Disease Control and Prevention. (June 4, 2010). Youth risk behavior surveillance-United States 2009. *Morbidity and Mortality Weekly Report: Surveillance Summaries, 59*(SS-5).
- 23. Centers for Disease Control and Prevention. (2007). Suicide trends among youths and young adults aged 10-24 years: United States 1990-2004. *Morbidity and Mortality Weekly Report, 56*(35), 905-908.
- 24. Dahlberg, L.L., Ikeda, R.M., & Kresnow, M. (2004). Guns in the home and risk of a violent death in the home: Findings

References continued

Information Dissemination in Schools

from a national study. *American Journal of Epidemiology, 160*(10), 929-936. doi:10.1093/aje/kwh309

- Brent, D.A., & Bridge, J.A. (2003). Firearms availability and suicide: Evidence, interventions, and future directions. *American Behavioral Scientist*, 46, 1192–1210. doi:10.1177/0002764202250662
- 26. Kalafat, J. (2003). School approaches to youth suicide prevention. *American Behavioral Scientist*, *46*(9), 1211-1223.
- Gould, M.S., Marrocco, F.A., Kleinman, M., Thomas, J.G., Mostkoff, K., Cote, J., & Davies, M. (2005). Evaluating iatrogenic risk of youth suicide screening programs: A randomized controlled trial. *JAMA*, 293(13), 1635–43.
- 28. Crawford, M.J., Thana, L., Methuen, C., Ghosh, P., Stanley, S.V., Ross, J, . . . Bajaj, P. (2011). Impact of screening for risk of suicide: Randomised controlled trial. *The British Journal of Psychiatry, 198,* 379-384. doi:10.1192/bjp.bp.110.083592
- 29. Hicks, B.B. (1990). *Youth suicide: A comprehensive manual for prevention and intervention*. Bloomington, IN: National Education Service.
- Moskos, M., Olson, L., Halbern, S., Keller, T., & Gray, D. (2005). Utah youth suicide study: Psychological autopsy. *Suicide and Life-Threatening Behavior*, 35(5), 536-546.
- Kaplan, S., Pelcovitz, D., Salzinger, S., Mandel, F., & Weiner, M. (1997). Adolescent physical abuse and suicide attempts. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36(6), 799-809.
- 31. Maine Center for Disease Control and Prevention. (2007). Maine Youth Suicide Prevention Implementation Plan.

Retrieved from http://www.state.me.us/suicide/myspp/ program/plan.htm

- 32. U.S. Public Health Service. (1999). *The Surgeon General's Call to Action to Prevent Suicide*. Washington, DC. Retrieved from http://www.surgeongeneral.gov/library/ calltoaction/
- Russell, S. T., & Joyner, K. (2001). Adolescent sexual orientation and suicide risk: Evidence from a national study. *American Public Health Association*, *91*(8), 1276-1281.
- 34. Bridge, J.A., Goldstein, T.R., & Brent, D.A. (2006) Adolescent suicide and suicidal behavior. *Journal of Child Psychology and Psychiatry*, *47*(3), 372-394.

Notes

Information Dissemination in Schools

Notes

Information Dissemination in Schools



Prepared by

Stephen Roggenbaum Justin Doan Katherine J. Lazear

Developed by

The Louis de la Parte Florida Mental Health Institute in the College of Behavioral and Community Sciences at the University of South Florida. Funded in part by the Orange County Department of Mental Health and Orange-Ulster BOCES. Originally funded by the Institute for Child Health Policy at Nova Southeastern University through a Florida Drug Free Communities Program Award.

Design & Page Layout Dawn Khalil © 2014, *Louis de la Parte* Florida Mental Health Institute



Stephen Roggenbaum roggenba@usf.edu 813-974-6149 (voice)



UNIVERSITY OF SOUTH FLORIDA COLLEGE OF BEHAVIORAL & COMMUNITY SCIENCES

Events, activities, programs and facilities of the University of South Florida are available to all without regard to race, color, marital status, gender, sexual orientation, religion, national origin, disability, age, Vietnam or disabled veteran status as provided by law and in accordance with the university's respect for personal dignity.

Permission to Copy all or portions of this publication is granted as long as this publication, the Department of Child & Family Studies, Louis de la Parte Florida Mental Health Institute, and the USF College of Behavioral & Community Sciences are acknowledged as the source in any reproduction, quotation or use.